REPORT OF TRANSFUSION-RELATED ACUTE LUNG INJURY (TRALI)

MEDIC TADI #:			
T	ime:		
rs after the start	of transfusion	on)	
ry distress Hypotension			
ry edema Fever (1-2°C rise)			
Obtain a serum sample from the components(s) and a pretransfusion sample from the patient.			
NOTIFY MEDIC HOSPITAL SERVICES IMMEDIATELY FOR RECALL OF OTHER COMPONENTS COLLECTED FROM SUSPECTED DONOR(S).			
		a	t MEDIC.
(Hospital Services Tech)			
By:			
BLOOD PRODUCTS TRANSFUSED 1-6 HOURS BEFORE ONSET OF SYMPTOMS:			
DATE TRANSFUSED	TIME STARTED	DATE COLLECTED	# PREVIOUS DONATIONS
DANGEUSION			TES
	Trs after the start	Time: rs after the start of transfusion Hypotension Fever (1-2 ⁴) s) and a pretransfusion sa EDIATELY FOR RECALL TED DONOR(S). ervices Tech) By: RS BEFORE ONSET OF S DATE TIME TRANSFUSED STARTED	Time: Time: rs after the start of transfusion) Hypotension Fever (1-2°C rise) s) and a pretransfusion sample from the EDIATELY FOR RECALL OF OTHER TED DONOR(S). a ervices Tech) By: a RS BEFORE ONSET OF SYMPTOMS DATE TIME DATE

INFORMATION COMLETED BY: _____DATE: _____DATE: _____